



## FERTILIZER W/ Barricade 0.45%

1702-0756

20-0-0

FOR SELECTIVE PRE-EMERGENCE CONTROL OF WEED GRASSES AND BROADLEAF WEEDS IN ALL ESTABLISHED TURF GRASSES, LAWNS, GOLF COURSE FAIRWAYS AND SOD FARMS, (NOT FOR PUTTING GREENS). USE FOR PRE-EMERGENCE WEED CONTROL AROUND LANDSCAPE ORNAMENTALS INCLUDING ESTABLISH PERENNIALS AND WILDFLOWER PLANTINGS.

### Active Ingredient:

Prodiamine\*\*\*..... 0.450%

Inert Ingredients:..... 99.550%

Total..... 100.000%

\*\*\*CAS 29091-21-2

EPA Reg. No. 52287-7

EPA Est. 52287-AL-001

### GUARANTEED ANALYSIS

Total Nitrogen (N)\*.....20.0000%

20.0000% Urea Nitrogen

Derived From: Polymer Coated Urea, Urea

\* 16% coated slow release Urea Nitrogen from Polymer Coated Urea

F352 Density - 70 lb./cu. ft.

Batch No. 1702-0756

Net Wt. 50 LBS. (22.7 KG)

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### READ ENTIRE LABEL BEFORE USING

#### FIRST AID

**IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, and then continue rinsing. Call a poison control center or doctor for treatment advice.

**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center or doctor, or when going for treatment. For information, call 1-800-282-8007.

### SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

##### ENVIRONMENTAL HAZARDS

This product has low solubility in water. At limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated area may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

##### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long sleeved shirt and long pants; chemical resistant gloves; shoes plus socks and a dust mask. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

##### USER SAFETY RECOMMENDATIONS

**Users should:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

##### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container away from other fertilizer, feed or food stuffs and separated from other pesticides. **PESTICIDE DISPOSAL:** If Empty: Do not reuse this container. Place in trash or offer for recycling if available. **IF Partly Filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**IF PRODUCT IS SPILLED:** Sweep up and use as per label instruction or dispose of as indicated above.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container, offer for recycling if available.

##### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This product is a turf fertilizer and a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings.

It controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 1/2 inch of rainfall, irrigation or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application. **Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farms, commercial nurseries).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required

for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls; chemical resistant gloves such as butyl rubber > 14 mils or nitrile rubber > 14 mils; shoes plus socks.

### NON-AGRICULTURAL USES REQUIREMENTS


The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The "WPS" applies when this product is used to produce agricultural plants on farms, forests or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry.

### APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment. Do not apply otherwise.

#### ESTABLISHED TURF

This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds listed below in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turfgrass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.



- 1: Mid-north & northwestern.....After May 30
- 2: Upper/mid-states & northeast.....After May 10
- 3: Lower mid-states.....After April 20
- 4: Upper southern states.....After March 20
- 5: Warm season states.....January 1 to March 20

#### USE PRECAUTIONS

The following precaution apply to the use of this product in turf grasses and lawns:

(1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as diking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

#### RATES OF APPLICATION

This product may be applied as a single application or in sequential application to control weeds germinating throughout the year. All applications must be made prior to germination of the weeds listed below. This product will not control established weeds. Maximum use rate section should be based on turf species. The length of time of residual pre-emergence weed control provided by this product is related to the rate applied.

## MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

### MAXIMUM APPLICATION RATE/CALENDAR YEAR OF PRODUCT BY TURF GRASS SPECIES<sup>(1)</sup>

Turf species	Lbs. product per acre	Lbs. A.I. Per acre
Creeping bentgrass	144 lbs.	0.65 lbs.
Creeping red fescue	167 lbs.	0.75 lbs.
Buffalograss	222 lbs.	1.00 lbs.
Kentucky bentgrass	"	"
Perennial ryegrass	"	"
Bermudagrass <sup>(2)</sup>	333 lbs.	1.50 lbs.
Bahiagrass	"	"
Centipedegrass	"	"
Seashore paspalum	"	"
St. Augustinegrass	"	"
Tall fescue (including turf type)	"	"
Zoysia	"	"

<sup>(1)</sup> These are the maximum rates per calendar year by species limitations. <sup>(2)</sup> May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 116 lbs. per acre. Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in springing situations.

Do not apply more than 333 lbs. of this product to the same area per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates section.

## WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barleygrass	Henbit	Purslane, Common
Bluegrass, Annual ( <i>Poa annua</i> )	Itchgrass	Pusley, Florida
Carp weed	Johnsongrass (from seed)	Rescuegrass <sup>(1)</sup>
Chickweed, Common	Junglerice	Shepherd's Purse <sup>(2)</sup>
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (Large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarter, Common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, Fall, Brown top)	Witchgrass
Goosegrass <sup>(1)</sup>	Pigweed	Woodsorrel, Yellow (from seed)

<sup>(1)</sup> In many areas a single application of 144 to 333 lbs. per acre will control Goosegrass. However, under heavy Goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 144 to 222 lbs per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates section. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates section. <sup>(2)</sup> Application for Shepherd's Purse should be made in late summer, fall or winter prior to germination. <sup>(3)</sup> Suppression only of Rescuegrass.

Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

## WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not that the phloem soil.

## WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and/or reseeding results.

Months Before Overseeding				
Product Rate Lbs./Acre	A.I./Acre	North	Transition	South
111	0.50	4	4	4
144	0.66	5	4	4
167	0.74	6	5	5
178	0.80	-	6	6
222	2.00	-	7	7
253	1.14	-	-	9
289	1.30	-	-	10
333	1.50	-	-	12

## APPLICATION EQUIPMENT AND SPREADER SETTINGS

The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application of this product. Follow directions in the owner's manual for calibrating your spreader.

## LANDSCAPE ORNAMENTALS INCLUDING ESTABLISHED WILD FLOWER PLANTINGS

This product may be applied for residual pre-emergence weed control in ornamentals listed on this label.

## USE RATES

Apply at 144 to 333 lbs. per acre (3.33 lbs. to 7.64 lbs. per 1,000 sq. ft.) of landscape plants, in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 333 lbs. per acre.

## APPLICATION TIMING AND INFORMATION

Apply uniformly to the surface of the soil in ornamental plantings beds with a granular applicator. Calibrate your applicator to assure appropriate dosage as described above under Use Rates, prior to application. Calibrate your applicator as described in the owner's manual. The bulk density of this product may vary; therefore, to assure that the product is applied at the rates described, you must use a calibrated applicator.

This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 1/4 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate post-emergence herbicide.

## ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

## NEWLY TRANSPLANTED ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplant and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. Irrigate after application as above directed.

## TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed here are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

## FOR USE AROUND LANDSCAPE ORNAMENTALS, INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWERS, AS LISTED.

Scientific Name	Common Name	Scientific Name	Common Name
Abies spp.	Fir Species** (Balsam, Fraser, Noble, Etc.)	Delosperma sp.	Cooperi Pink
Abella grandiflora	Abelia: Sherwood	Delphinium sp.	Larkspur: Blue Elf
Acer palmatum	Japanese Maple	Dianthus deltoids	Dianthus: Maiden Pink "Zing"
A. platnoides	Norway Maple	D. graziopolitans	Cheddar, Pink
Achillea sp.	Yarrow: King Edward	Dodonea viscosa	Hop Bush
Actinidia chinensis*	Kiwi*	Echinacea purpurea	Coneflower, Purple
Agapanthus africanus	Lily-of-the-Nile (African Lily)	Elaeagnus pungens	Magnus Silverberry
Agapanthus orientalis		Euonymus fortunei	Wintercreeper
Akebia quinata	Five-Leaf or Chocolate Vine	E. japonica	Japanese Spindle Tree (Evergreen Euonymus)
Allium cernuum	Lady's Leek, Nodding	E. kiautschovica	Spreading Euonymus
Anemone hybrida		Fatsia japonica	Japanese Aralia
Aquilegia sp.	Japanese Anemone	Forsythia intermedia	Border Forsythia
Arctostaphylos densiflora	Vine Hill Manzanita	F. suspense	Weeping Forsythia
Arctotheca calendula	Cape Weed	F. viridissima	Greenstem Forsythia
Artemisia sp.	Wormwood: Silver Mound, Castle	Gaillardia sp.	Gaillardia, Blanket Flower: "Goblin"
Aster sp.	Aster: Bonny Blue, Purple Dome	Gardenia jasminoides	Gardenia, Cape-Jasmine, Radicans

Aster X Frikartii	Gaua sp.	Gentian
Athrium Filiz-femina	Lady Fern: Fern Lady	Gentiana dabrura
Aucuba japonica	Japanese Aucuba	Gernium cinereum
Begonia sp.	Fibrous Begonia:	Gladiolus spp.**
	Hardy Grandis	
Berberis gladwynensis	Barberry	
B. julianae	Wintergreen	Gypsophila repens
B. mentorensis	Barberry	Hedera helix
B. thunbergii	Mentor Barberry	Helianthemum sp.
	Japanese Barberry	Heimer callis sp.
B. verruculosa	Warty Barberry	Hibiscus**
Bergenia cordifolia		Hibiscus sp.
Boltonia asteroides	Snowbank	Hibiscus rosa-sinensis
Bougainvillea spp.	Bougainvillea	Hosta plantaginea
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red	H. Sieboldiana Hosta
Scientific Name	Common Name	Scientific Name
Buxus microphylla	Japanese Boxwood	Houttuynia cordata var. variegata
Callistemon viminalis	Weeping Bottlebrush	Hydrangea macrophylla
Calhna vulgaris	Scotch Heather	Lex cornuta
Campanula carpatica	Tussock Bellflower (White Clips)	L. crenata
Campis X Tagliabuana	Trumpet Creeper, Trumpet Flower, Madame Galen	I. opaca
Capobrotus edulis	Hotentot Fig (Ice Plant)	I. pernyi
Cassia artemisoides	Feathery Cassia	I. vomitoria Yaupon
Ceanothus rigidus	Wild Lilac	lnula ensifolia
Ceratostigma		Iris ensata
Plumbago noides		Sword-Leaved Iris; Jodlesong
Chamaecyparis		
Pisifera	False Cypress	Iris siberica
Chrysanthemum nipoicum		Iris spp.**
Cleyera japonica	Cleyera	Jasminium udidorum
Citrus spp.*	Citrus species*	Juniperus chinensis
Coreopsis sp.	Coreopsis (Calliopsis): J. conferta	Winter Jasmine
	Early Sunrise, Moonbeam	Chinese Juniper; Nick's Compact, Parsonii
Cornus florida	Flowering Dogwood	Shore Juniper; Blue Pacific
C. stolonifera	American Dogwood	J. davurca
Cortaderia selloana	Pampas Grass	J. horizontalis
Contoneaster apiculatus	Cranberry	Juglans sp.*
	Cotoneaster	Justicia brandegeana
C. buxifolius	Cotoneaster	Lagerstromia
		Shrimp Plant
C. dammeri	Barberry	Indica Lagerstromia
	Cotoneaster	Indica X Fauriei
	Rockspray	montevidenensis
C. microphyllus	Cotoneaster	Lavender sp.
	Hawthorne	Lavender; Munstead
Cretaeus sp.		Leontopodium alpinum
		Edelweiss
Crococsmia sp.	Lucifer	Ligustrum sinense
		Chinese Privet; Variegata
Cupressus sempervirens	Italian Cypress	L. amurense
Delosperma alba	White Trailing Ice Plant	L. japonicum
	Glossy Privet (Waxleaf)	Veronica sp.
L. juicidum		Veronica, Speedwell; Sunny Border
Lilium sp.	Lily; Jazz Big Blue Lily; Turf	Viburnum japonicum
		Japanese Viburnum
Liriope muscari	Silver Mound; Evergreen Giant	V. odoratissimum
	Liriope, Variegated	Sweet Viburnum
Liriope Muscari var. variegata		V. plicatum
Liriope spicata	Liriope, Creeping	V. rigidum
		Japanese Snowball
Lobelia Cardinalis	Cardinal Flower, Indian Pink	V. suspensum
		Camary Island Viburnum
Lonicera japonica	Japanese Honeysuckle	V. timus
		Arrow wood Viburnum
		Laurustinus

L. tatarica	Tatarian Honey-suckle	V. trilobium	Cranberry Bush
Loropetalum chinense	Burgundy	V. Wrightii	Leatherleaf Viburnum
Lythrum sp.	Looseleaf, Modern Pink	Vinca major	Greater Periwinkle,
Magnolia spp.**	Magnolia species**	Vinca minor	Vinca
Maleophora huteola	Ice Plant	Vitis sp.*	Common Periwinkle,
Malus sp.*	Crabapple*	Weigela florida	Vinca
Miscanthus sinensis	Yaku Jima, Silberfeder	Yucca aloifolia	Grape*
Nandina domestica	Heavenly Bamboo	Y. filamentosa	Old Fashioned Weigela
Narcissus spp.**	Narcissus species**		Spanish Bayonet,
Nerium oleander europea*	Oleander Olea Olive*		Yucca
Ophiopogon japonicus	Mondo Grass		Adam's Needle,
Osmanthus heterophyllus	Osmanthus (False Holly); Gulf Tide		Yucca
Osteospermum fruticosum	Trailing African Daisy		
Oxydendrum arborescens	Sourwood		
Paeonia suffruticosa	Tree Peony		
Pennisetum setaceum	Fountain Grass (Dwarf)		
Perovskia atriplicifolia	Avocado*		
Persea americana	Frasier's Photinia (Redtip)		
Photinia fraseri	Dragonhead, False; Vivid		
Physotegia Virginiana	Spruce species** (Colorado)		
Picea spp.**	Blue, Norway, etc.)		
Pieris japonica	Lily-of-the-Valley Bush		
Pinus brutia	Calabrian Pine		
P. canariensis	Canary Island Pine		
P. eliotii	Splash Pine		
P. halepensis	Aleppo Pine		
P. nigra	Australian Pine		
P. palustris	Longleaf Pine		
P. radiata	Monterey Pine		
P. thunbergiana	Japanese Black Pine		
P. strobus	Eastern White Pine		
P. sylvestris	Scotch Pine		
P. taeda	Loblolly Pine		
P. virginiana	Virginia Pine		
Pistachio sp.*	Pistachio*		
Pittosporum rhombifolium	Queensland Pittosporum		
P. tobita	Japanese Pittosporum		
Podocarpus macrophyllus	Japanese Yew		
Prunus faureroensis	English Laurel		
Prunus sp.*	Almond, Apricot, Nectarine, Peach, Plum and Prune*		
Pseudotsuga menziesii**	Douglas Fir**		
Pyracantha coccinea	Firethorn (Scarlet)		
P. fortuneana	Firethorn		
P. koidzumii	Firethorn		
Pyrus sp.	Bradford Pear sp.		
Quercus shumardii	Oak, Shumard's Red		
Quercus spp.	Oak, species		
Raphiolepis indica	Indian Hawthorn		
Raphiolepis umbellata	Yedda Hawthorn		
Rhododendron 'Coral Bells' (including Azalea)	Delaware Valley White		
	'Flame Creeper'		
	'Formosa'		
	'Girard Crimson'		
	'George L. Tabor'		
	'Hino-crimson'		
	'PJM'		
	'Roseum Elegans'		
	'Wakefield'		
	'White Gumbo'		

Rosa banksiae	Lady's Bank's Rose
Rosmarinus officinalis*	Rosemary*
Rudbeckia sp.	Black-Eyed Susan;
	Goldstrum
Santolina, virens	Santolina
Scientific Name	Common Name
Saxifraga sp.	Saxifrage; Purple
	Dome Pincushion
	Flower
Scabiosa sp.	Stonecrop
Sedum canticola	Stonecrop; Lidakense
Sedum dasyphyllum	Stonecrop Sedum
spuriatum	Stonecrop; Dragon's
	Blood
Spiraea Bumalda	Spiraea; Anthony
	Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew

\* Do not use on food producing trees, vines, or plants \*\* Not for use on container grown plants

### CONDITIONS OF SALE AND WARRANTY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Harrell's, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Harrell's, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Harrell's, LLC disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

**LIMITATION OF WARRANTIES:** The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Harrell's, LLC option, the replacement of the product.

**Harrell's, LLC**  
**P. O. Box 807**  
**Lakeland, FL 33802**

(F352)

Barricade® is a registered trademark of Syngenta, Inc.